This listing of claims will replace all prior versions, and listings, of claims in the application:

In the Claims:

1. (Previously Presented) A process for qualitative and/or quantitative detection of analytes in a liquid and/or solid phase homogeneous immunoassay, comprising determining remanence magnetization in said homogeneous immunoassay of a stable of quasi-stable ferromagnetic or ferrimagnetic substance bound to said analyte, wherein said homogeneous immunoassay has been contacted with said ferromagnetic substance, whereby said substance binds to the analyte, and the presence of such magnetization is indicative of the presence or amount of the analyte.

(Previously Presented) A process for qualitative and/or quantitative detection of analytes in homogeneous immunoassays comprising measuring remenance magnetization of magnetic markers in a sample, bound to said analytes, wherein said homogenous immunoassay has been contacted with said ferromagnetic substance, whereby said substance binds to the analyte, and at the time of measurement the magnetization of unbound magnetic markers that are present in the sample in their totality fades owing to extrinsic superparamagnetism, and the presence of remenance magnetization is indicative of the presence or amount of said analytes.

- 27

 2. (Previously Presented) A process for qualitative and/or quantitative detection of analytes in a liquid or solid phase homogeneous immunoassay, comprising
 - (i) labeling first structure-specific substances, with ferrimagnetic or ferromagnetic substances to indicate the presence or amount of said analyte,
 - (ii) adding said magnetic labeled structure-specific substances to a sample that is to be measured.
 - (iii) magnetizing the sample to be measured with the aid of a magnetic field or suitable intensity that is applied from outside and,
 - (iv) measuring the remanence of the magnetization of bound structure-specific substances with the sid of magnetic field sensors after the external field is shut off, without removing unbound structure-specific substances.

(Canceled) 22

(Previously Presented) The process according to claim 46; wherein the structure-specific substances are antibodies, antibody fragments, bioun, substances that bind specifically to biotic, agomsts that bind specifically to receptors of their antigonists, peptides, proteins, receptors, enzyme substrates, nucleotides, ribonucleic, acids, deoxyribonucleic acids, carbohydrates, or lipoproteins.

28

8. (Previously Presented) The process according to claim 3, wherein the structure-specific substances have a binding constant in the range of 10³-10¹⁵ (mol/l)⁻¹.

29
27
2. (Previously Presented) The process according to claims, wherein the structure-specific substances have a binding constant in the range of 10⁷-10¹⁵ (moVi)⁻¹.

2. (Previously Presented) The process according to claim 1, wherein the sample is moved during the measurement and a sample signal is modulated.

3

87. (Previously Presented) The process according to claim 1, wherein induction coils that are hooked up as gradiometers, flux gate-magnetometers, giant magnetoresistance sensors, or magnetoresistaive converters are used as magnetic field sensors to determine remanent magnetization.

40. (Previously Presented) The process according to claim 1, wherein SQUIDs are used as magnetic field sensors to determine remanent magnetization.

(Previously Presented) The process according to claim 1, wherein simultaneous determination of several different analytes in a sample of liquids or solid substances is carried out by sequential magnetization of a sample to be measured.

6

M. (Previously Presented) The process according to claim M, wherein for simultaneous quantitative determination of analytes, different ferromagnetic or ferrimagnetic substances with discrete coercive field intensities are used.

- (Previously Presented) The process according to claim 1, wherein the ferromagnetic and ferrimagnetic substances used have intrinsic Neelian relaxation times greater than the measuring time.
- 9
 14. (Previously Presented) The process according to claim 15, wherein the ferromagnetic and ferrimagnetic substances that are used have Neelian relaxation times longer than 10° seconds at 20°C.

16. (Previously Presented) The process according to claim 16, wherein the ferromagnetic and ferrimagnetic substances that are used have Neelian relaxation nimes longer than 1 second at 20°C.

(Previously Presented) The process according to claim 1, wherein the ferromagnetic and ferrimagnetic substances have a particle size of 1 to 1000 nm.

1. (Previously Presented) The process according to claim 1, wherein the ferromagnetic and ferrimagnetic substances have a particle size in the range of 2 to 500 mm.

13
16. (Previously Presented) The process according to claim 1, wherein the ferromagnetic and ferrimagnetic substances are stabilized with a shell of oligomeric or polymeric carbohydrates, proteins, peptides, nucleotides, surfactants, synthetic polymers, and/or lipids.

19. Canceled

12

- 20. Canceled
- 21. Canceled

20 27

2. (Previously Presented) The process according to claim, wherein the structure-specific substances are cytokines, lymphokines, endothelins or their antagonists.

14

- 23. (Previously Presented) The process according to claim 1, wherein the ferromagnetic or ferrimagnetic substances are stable or quasi-stable colloidal particles that are made of iron oxides, barium ferrise, strontium ferrite, pure iron, chromium, dioxide, nickel, and cobalt, or iron oxides with manganese, copper, nickel, or cobalt additives.
- 7
 24. (Previously Presented) The process according to claim 17. wherein several ferromagnetic or ferrimagnetic substances with various coercive field intensities are used.
- 25. (Previously Presented) in a fertility, histocompatibility, allergology, infectiology, hygiene, genetics, virology, bacteriology, toxicology, pathology, environmental analysis, or medical diagnosis process comprising detecting an analyte; the improvement wherein the detecting is performed according to claim 1.

16

- 26. (Previously Presented) The process according to claim 1, wherein ferromagnetic or ferromagnetic substances are introduced into the human body or are applied on the human body, and the remanence of the magnetization of the ferromagnetic of ferromagnetic substances is determined after a magnetizing field is shut off.
 - 3₁ 27
- 27. (Previously Presented) The process according to claim 3/, wherein ferromagnetic or ferromagnetic substances are introduced into an organism or applied on the organism, by a process comprising
 - labeling structure-specific substances with terromagnetic or ferromagnetic substances,
 - (ii) adding said magnetic labeled structure-specific substances to a living organism or applied to an organism,

- (III) magnetizing a volume of the organism with the aid of a magnetic field that is applied from the outside and,
- (iv) measuring remanence of the magnetic markers with the aid of magnetic field sensors after the external field is shut off.

22. (Previously Presented) The process according to claim 27, wherein antibodies, antibody fragments, agonists that bind specifically to receptors or their antagonists, peptides, proteins, receptors, enzymes, enzyme substrates, nucleotides, ribonucleic acids, deoxyribonucleic acids, carbohydrates, or lipoproteins are used as structure-specific substances.

32.
35. (Previously Presented) The process according to claim 26, wherein the agonists or antagonists that bind specifically to receptors are cytokines, lymphokines, endothelines or their antagonists.

32.
30. (Previously Presented) The process according to claim 28, wherein the structure-specific substances have a binding constant in the range of 10⁵-10¹⁵ (mol/1)⁻¹.

35
36
37. (Previously Presented) The process according to claim 28; wherein the structure-specific substances have a binding constant in the range of 10⁷-10¹⁵ (moV1).

(Previously Presented) The process according to claim 26, wherein Superconducting Quantum Interference Devices (SQUIDs), induction coils, fluxgate-magnetometers, giant magnetoresistance sensors, or magnetoresistive converters are used as magnetic field sensors.

- 33. Canceled
- 34. Canceled

21

3/5. (Previously Presented) The process according to claim 77, wherein a mixture of different ferrimagnetic or ferromagnetic substances with structure-specific substances is used.

18
26. (Previously Presented) The process according to claim 26, wherein the Neclian relaxation time of the ferromagnetic or ferrimagnetic substances is longer than 10⁻⁶ second at 37°C.

M. (Previously Presented) The process according to claim 26, wherein the Neclian relaxation time of the ferromagnetic or ferrimagnetic substances is longer than 1 second at 37°C.

18
38. (Previously Presented) The process according to claim 36, wherein the ferrimagnetic or ferromagnetic substances are iron oxides or iron oxides with manganese, copper, nickel, or cobalt additives.

49. (Previously Presented) The process according to claim 1, wherein the ferromagnetic or ferromagnetic substance is magnetic-labeled anticollagen II and SQUID(s) are used to determine remanent magnetization.

40. (Previously Presented) A process according to claim 1, wherein the ferromagnetic or ferrimagnetic substances are used to label structure specific substances, which are added to the analyte.

A1. (Previously Presented) A process according to claim 1, wherein the analyte is labeled with structure specific substances, and the ferromagnetic or ferrimagnetic substances are added thereto.

26
42. (Previously Presented) A process according to claim at, wherein the structure-specific substances are antibodies, antibody fragments, biodin, substances that bind specifically to biotin, agonists that bind specifically to receptors of their antagonists, peptides, proteins, receptors, enzyme substrates, nucleotides, ribonucleic, acids, deoxyribonucleic acids, carbohydrates, or lipoproteins.